

Letter to the editor

Note on the cubic decreasing region of the Chebyshev method

T. Zhanlav

National University of Mongolia, Department of Applied Mathematics, PO 46/687 Ulaanbaatar 210646, Mongolia

ARTICLE INFO

Article history:

Received 28 October 2009

Received in revised form 21 May 2010

MSC:

47H17

65J15

Keywords:

Nonlinear equations on Banach spaces

Chebyshev method

Region of convergence

ABSTRACT

In this paper we reduce the two-dimensional cubic decreasing region considered in Hernandez and Salanova (2000) [1], [2] into one-dimensional region or interval for the Chebyshev method. It means that we find a simple sufficient condition for the semilocal convergence of the method.

© 2010 Elsevier B.V. All rights reserved.

1. Introduction

We consider a nonlinear equation

$$F(x) = 0. \quad (1)$$

Here $F : \Omega \subseteq X \rightarrow Y$ is a nonlinear, twice Frechet differentiable operator defined on a convex nonempty domain Ω and X and Y are Banach spaces. Chebyshev method is one of the best known third-order iterative process for solving the nonlinear equation (1) and it is given by

$$\begin{aligned} y_n &= x_n - \Gamma_n F(x_n), \quad \Gamma_n = F'(x_n)^{-1}, \\ x_{n+1} &= y_n - \frac{1}{2} \Gamma_n F''(x_n)(y_n - x_n)^2, \quad n = 0, 1, \dots \end{aligned} \quad (2)$$

Let us assume that $F'(x_0)^{-1} \in \mathcal{L}(Y, X)$ exists for some $x_0 \in \Omega$, where $\mathcal{L}(Y, X)$ is the set of bounded linear operators from Y into X .

Moreover, we suppose that [1,2]

- (c₁) $\|\Gamma_0\| \leq \beta$,
- (c₂) $\|y_0 - x_0\| = \|\Gamma_0 F(x_0)\| \leq \eta$,
- (c₃) $\|F''(x)\| \leq M, \quad x \in \Omega$,
- (c₄) $\|F''(x) - F''(y)\| \leq K\|x - y\|, \quad x, y \in \Omega, \quad K > 0$

and

$$a_0 = M\beta\eta > 0, \quad b_0 = K\beta\eta^2 > 0. \quad (3)$$

E-mail address: zhanlav@yahoo.com.




Fig. 1. Cubic decreasing regions.

In [1,2] the semilocal convergence theorem for the Chebyshev method was proven under the conditions (c_1) – (c_4) and

$$0 < a_0 < 1/2, \quad b_0 < \frac{3}{4}(1 - 2a_0)(a_0 + 2)(4 - 2a_0 - a_0^2). \quad (4)$$

The cubic decreasing regions [1] of the Chebyshev method are presented in Fig. 1, where a_0 and b_0 are taken as coordinates.

The dotted line represents the curve

$$b_0 = \frac{6(1 - 2a_0)(4 - 2a_0 - a_0^2)}{(2 + a_0)^2}$$

obtained in [3] and the continuous line represents the curve

$$b_0 < \frac{3}{4}(1 - 2a_0)(a_0 + 2)(4 - 2a_0 - a_0^2)$$

obtained in [1].

2. The cubic decreasing region

It is evident that the cubic decreasing region given in [1] is bigger, and therefore, the region of accessibility of the Chebyshev method has been increased. However the second inequality in (4) is not satisfied for $0 < a_0 < 1/2$ and it may hold only for the region $0 < a_0 < a_0^*$, where $a_0^* < 1/2$. Therefore, it is desirable to obtain a one-dimensional cubic decreasing region $0 < a_0 < a_0^*$ instead of the two-dimensional one given in (4) for the Chebyshev method. To this end, we proceed as follows: From (3) it is clear that there exists a closed relationship between b_0 and a_0 . More precisely, we have

$$b_0 = c_n a_0^n, \quad \text{with } c_n = \frac{K}{M^n \beta^{n-1} \eta^{n-2}}, \quad n = 1, 2, \dots$$

When $n > 2$, the coefficient c_n will increase due to the small η and as a consequence, the cubic decreasing region will be reduced. From this, it is clear that it suffices to consider only the cases $n = 1$ and $n = 2$.

We first consider the case $n = 1$, i.e.

$$b_0 = c_1 a_0, \quad (5)$$

where

$$c_1 = \frac{K\eta}{M} > 0. \quad (6)$$

The second inequality in (4) leads as

$$\psi_1(a_0) = 2a_0^4 + 7a_0^3 - 4a_0^2 - \left(16 + \frac{4}{3}c_1\right)a_0 + 8 > 0. \quad (7)$$

Since $\psi_1(0) = 8 > 0$ and $\psi_1(1/2) = -\frac{2}{3}c_1 < 0$, there exists one value a_0^* , such that $\psi_1(a_0^*) = 0$ ($a_0^* < 1/2$). Differentiating (7) with respect to a_0 we obtain

$$\psi_1'(a_0) = 8a_0^3 + 21a_0^2 - 8a_0 - \left(16 + \frac{4}{3}c_1\right).$$

From this, it follows that $\psi_1'(a_0) < 0$ under $0 < a_0 < 1/2$, i.e. $\psi_1(a_0)$ is a decreasing function on the interval $0 < a_0 < 1/2$ and therefore, the inequality (7) holds for $0 < a_0 < a_0^*$.




Fig. 2. The convergence region, $n = 1$ case.

We cannot find the point a_0^* , but we find the upper sharp bound for it:

$$a_0^* \leq \hat{a}_0 = \frac{1}{2 + c_1}.$$

Indeed, we have

$$\psi_1(\hat{a}_0) = \frac{c_1}{3z^4}(20z^3 - 12z - 3),$$

where $z = c_1 + 2$ and $20z^3 - 12z - 3 > 0$ for $2 \leq z < \infty$. This means that

$$\psi_1(\hat{a}_0) > 0, \quad \text{when } c_1 > 0$$

$$\psi_1(\hat{a}_0) = 0, \quad \text{when } c_1 = 0.$$

Thus the one-dimensional (cubic decreasing) region for the Chebyshev method is

$$0 < a_0 < a_0^*, \quad (8)$$

where a_0^* is a root of the equation $\psi_1(a_0) = 0$, lying in $(0, 1/2)$, as shown in Fig. 2.

Now we consider the case of $n = 2$, i.e.

$$b_0 = c_2 a_0^2, \quad c_2 = \frac{K}{M^2 \beta} > 0.$$

In this case the second inequality in (4) leads to

$$\psi_2(a_0) = 2a_0^4 + 7a_0^3 - 4\left(1 + \frac{c_2}{3}\right)a_0^2 - 16a_0 + 8 > 0. \quad (9)$$

Since $\psi_2(0) = 8 > 0$, $\psi_2(1/2) = -\frac{1}{3}c_2 < 0$ and

$$\psi_2'(a_0) = 8a_0^3 + 21a_0^2 - 8\left(1 + \frac{c_2}{3}\right)a_0 - 16 < 0 \quad \text{for } 0 < a_0 < 1/2$$

then $\psi_2(a_0)$ is a decreasing function with respect to a_0 and therefore, there exists $\bar{a}_0 < 1/2$ such that $\psi_2(\bar{a}_0) = 0$. It means that $\psi_2(a_0) > 0$ when $0 < a_0 < \bar{a}_0$. As before, we can only find the upper bound for \bar{a}_0 :

$$\bar{a}_0 \leq \tilde{a}_0 = \frac{1}{2 + c_2}.$$

Simple calculation gives

$$\psi_2(\tilde{a}_0) = \frac{1}{z^4} \left\{ 8z^4 - 16z^3 - 4\left(1 + \frac{c_2}{3}\right)z^2 + 7z + 2 \right\} = \frac{z-2}{3z^4} (24z^3 - 4z^2 - 12z - 3),$$

where $z = 2 + c_2 \geq 2$.

The function $24z^3 - 4z^2 - 12z - 3$ is an increasing function of $z \in [2; +\infty)$. Therefore, we have

$$\psi_2(\tilde{a}_0) > 0, \quad \text{when } c_2 > 0$$

$$\psi_2(\tilde{a}_0) = 0, \quad \text{when } c_2 = 0.$$




Fig. 3. The convergence region, $n = 2$ case.

Thus the inequality (4) holds under condition

$$0 < a_0 < \tilde{a}_0, \quad (10)$$

where \tilde{a}_0 is root of equation $\psi_2(a_0) = 0$, lying in $(0; 1/2)$, as shown in Fig. 3.

From (4), (9) and (10) we can conclude that the semilocal cubic convergence of the Chebyshev method holds under the conditions (c_1) – (c_4) and

$$0 < a_0 < \max\{\hat{a}_0, \tilde{a}_0\}.$$

Acknowledgements

The author is grateful to the anonymous referee for very important suggestions and comments to improve the first version of the paper. The research reported herein was sponsored in part by the foundation of Science and technology of Mongolia.

References

- [1] M.A. Hernandez, M.A. Salanova, Modification of the Kantorovich assumptions for semilocal convergence for the Chebyshev method, *J. Comput. Appl. Math* 126 (2000) 131–143.
- [2] M.A. Hernandez, Second-derivative-free variant of the Chebyshev method for nonlinear equations, *J. Optim. Theory Applications* 104 (2000) 501–515.
- [3] V. Candela, A. Marquina, Recurrence relations for rational cubic methods II: the Chebyshev method, *Computing* 45 (1990) 355–367.